When to get your COVID-19 vaccination



PRIMARY DOSES

	PFIZER	MODERNA	NOVAVAX	ASTRAZENECA*
	5 years old and over	6 years old and over	18 years old and over	18 years old and over
Interval between Primary dose 1 and 2	8 weeks Can be reduced to 3 weeks in some circumstances	8 weeks Can be reduced to 4 weeks in some circumstances	3 weeks Can be extended to 8 weeks in some circumstances	12 weeks Can be reduced to 4 weeks in some circumstances
Interval between Primary dose 2 and 3 for people who are severely immuno-compromised	2 months	2 months	2 months	2 months Not preferred as a third dose but can be used if there is no alternative

^{*} The Pfizer, Moderna or Novavax COVID-19 vaccines are preferred over the AstraZeneca vaccine for people under 60 years of age. However, adults under 60 years of age may still choose to receive the AstraZeneca vaccine if they have made an informed decision after considering the risks and benefits. Talk to your doctor to help make your decision.

If you have tested positive to COVID-19, it is recommended that you wait 3 months from your positive test result for your next COVID-19 vaccine dose.

For more information, visit www.health.gov.au or call the National Coronavirus Helpline on 1800 020 080. Select option 8 for free interpreting services.



When to get your COVID-19 vaccination







BOOSTER DOSE

for everyone aged 12 years

Booster dose

and older

PFIZER	MODERNA	NOVAVAX	ASTRAZENECA	
12 - 15 years old	18 years old and over			
3 months after primary doses	3 months after primary doses	3 months after primary doses	3 months after primary doses	
Only recommended for people aged 12 to 15 years old who:		Can be used if no other COVID-19 vaccine brand is suitable	Not preferred as a booster dose but can be used if no other COVID-19 vaccine	
 are severely immunocompromised 			brand is suitable	
 have a disability with significant or complex health needs 				
 have complex and/or multiple health conditions that increase the risk of severe COVID-19. 				
16 years and over				
3 months after primary doses				

If you have tested positive to COVID-19, it is recommended that you wait 3 months from your positive test result for your next COVID-19 vaccine dose.

For more information, visit www.health.gov.au or call the National Coronavirus Helpline on 1800 020 080. Select option 8 for free interpreting services.



When to get your COVID-19 vaccination



FOURTH BOCE					
FOURTH DOSE	PFIZER	MODERNA	NOVAVAX	ASTRAZENECA	
A fourth COVID-19 vaccine dose for people who are:	3 months after	3 months after	3 months	3 months	
 50 years and older 	booster dose	booster dose	after primary doses	after primary doses	
 residents of aged care or disability care facilities 			Can be used if no	Not preferred as a	
 16 years and older if they: 			other COVID-19	winter dose but can	
			vaccine brand is suitable	be used if no other COVID-19 vaccine	
are severely immunocompromisedhave, or have survived, certain cancers			Suitable	brand is suitable	
 have an autoimmune disease requiring treatment 					
 have a chronic or severe disease that affects their lungs, liver, kidneys, brain or heart 					
- have diabetes requiring medication					
 have a disability with significant or complex health needs 					
- are severely overweight or underweight.					
This will be a fifth COVID-19 vaccine dose for people who are severely immunocompromised.					
People aged 30 to 49 years can also choose to have a fourth COVID-19 dose.					
Speak with your doctor if you are not sure if you should receive a fourth COVID-19 dose.					

If you have tested positive to COVID-19, it is recommended that you wait 3 months from your positive test result for your next COVID-19 vaccine dose.

For more information, visit www.health.gov.au or call the National Coronavirus Helpline on 1800 020 080. Select option 8 for free interpreting services.

